Inform us about your routine medications and any diseases.

Do not eat or drink

before your surgery.

anything, including water, after midnight the night

The night before surgery, sho

Wax or shave your bikini area

and legs. It may be uncomfort

wer and wash the surgical

areas with hygienic soap.

able to do so in the days

Wear only comfortable, loose-fitting clothing that

button or zips in the

<u>info@longevita.co.uk</u> ର longevita.co.uk


front.

immediately after surgery.

Pre-operative



Do not take Aspirin, Alieve, or







Ibuprofen. Review the list of drugs containing aspirin and ibuprofen carefully.




/longevitauk





Stop smoking for 2 weeks before the surgery, and avoid alcohol consumption one week prior to the surgery.







Do not wear make-up, moisturizers, creams, lotions on the day your procedure is scheduled.



Do not wear jewelry, contact lenses and remove any piercings on the day of the procedure.




/c/longevita



AT to EXPECT?

Post-Operative

Return to hotel. Expect some swelling and bruising, mild discomfort.



swelling.


Expect maximum bruising and

You are about to invest time and money in your treatment abroad. Please follow these instructions carefully. If you have any queries, please do not hesitate to contact your Travel Coordinator or email to <u>info@longevita.co.uk</u>

Itching

6 longevitauk

Post-Operative



residual damaged fat cells. For every fat

You are about to invest time and money in your treatment abroad. Please follow these instructions carefully. If you have any queries, please do not hesitate to contact your Travel Coordinator or email to <u>info@longevita.co.uk</u> 6 longevitauk /longevitauk

or lie down until it passes. Like with any surgical procedure, you can expect some pain and discomfort during your initial recovery period as your tissue

/longevitauk

begins to heal. There is a very good chance that you have extensive bruising and swelling in the areas where the fat cells were taken from, and in the area where they were injected. This pain can be managed with prescribed painkillers.

/c/longevita

Patients may experience a brief sensation

Feeling lightheaded is similar to what you

might experience when standing-up too quickly. Should dizziness occur, simply sit

of dizziness the morning after surgery.

Expected Discomfort

VEAT to EXPECT? Numbress Depression You may feel numbness in your surgical

It's normal to have swelling in the surgical area for about six weeks. In some cases, swelling might take up to three months to resolve. You are about to invest time and money in your treatment abroad. Please follow these instructions carefully. If you have any queries, please do not hesitate to contact your Travel Coordinator or email to

/longevitauk

/longevitauk /c/longevita

Health, Beauty, and Wellness
Personel Care

Activities

It is normal to feel a bit anxious or

following surgery; but generally, after

about a week, you should begin to feel

depressed in the days or weeks

better.

You shouldn't do any heavy exercise or strenuous activity for at least a month after the procedure. Please avoid drinking alcohol while using the pain medications. After a full week of recovery, walking is greatly encouraged. Walking will improve your blood circulation, speed your healing, and might even improve your results!

Health, Beauty, and Wellness
Personel Care Post-Operative

You may start driving 2-3 weeks after the surgery. Please discuss it with your patient relations specialist. Eat Healty

You need to conform to a high fat, high calorie diet for 3 months

following the BBL procedure. This type of diet is necessary in order to maintain the fullness and volume of your buttocks. Eating a diet high in fat and calories is also vital in providing nourishment to the newly transferred

Post-Operative

• It is really important do exercise once you have completed your first month. in your treatment abroad. Please follow these instructions carefully. If you have any queries, please do not hesitate to contact your Travel Coordinator or email to

Do not drive for three weeks.

• Do not swim for four weeks.

1 month.

/longevitauk

Do not engage in sexual intercourse for

/c/longevita

Sitting Sleeping

circulation which could injure the fat grafts before they incorporate and become stable. Do not sit, bend over, or lay on your back until cleared by your surgeon.

Instead, you should remain standing or

rest by lying on their sides or stomach.

If you absolutely must sit, it is adviced

/c/longevita

to use a pillow for your comfort.

area of the fat graft will limit blood

Most people return to work within two to damage.

your final results.

Discuss returning to work with your patient

Take Care! You are about to invest time and money in your treatment abroad. Please follow these instructions carefully. If you have any queries, please do not hesitate to contact your Travel Coordinator or email to

Much of swelling has begun to go away. (once you have completed your first month.) You can remove your tape and take a shower. (f) /longevitauk /longevitauk /c/longevita Expected Discomfort

Itching of the treated areas several days after surgery may occur as part of the normal healing process. Small Bumps

cell that we remove during liposuction, we injure a few more. This is a beneficial effect because your body will be healing, gradually.

areas for months or even years. It's normal to have bruises as well. You may have fluid-filled swelling

will eventually fade.

above the scar, but this will go away.

Your scar may be red and raised, but it

Post-Operative

Inflammation

Resulting

7260

Post-Uperative

s and

<u>info@longevita.co.uk</u>

⊖ longevitauk

If you are a smoker, you should not smoke for at least 2 weeks after surgery.

Smoking and chewing tobacco

inhibit your circulation and can

significantly compromise your

You are about to invest time and money

(f) /longevitauk

surgical outcome.

info@longevitá.co.uk

6 longevitauk

Keep access sites clean, dry and inspect daily for signs of infection.

Minimize sun exposure to the

surgical areas for at least 12

Always use a sunscreen with a high SPF when going out in

months.

the sun.

fat cells. You are about to invest time and money in your treatment abroad. Please follow these instructions carefully. If you have any queries, please do not hesitate to contact your Travel Coordinator or email to <u>info@longevita.co.uk</u> 6 lonvegitauk (II) /longevitauk

Return to work varies with your type of

work, your overall health and personal

four weeks.

preferences.

care specialist.

<u>info@longevita.co.uk</u>

Slongevitauk

/longevitauk

When Can I Get Back to Work?

Be Patient for A Great Result!

The most important aspect of recovering from the procedure is the avoidance of pressure on the buttocks before it's fully healed. Any pressure applied to the recipient

/longevitauk

Be Patient!

/longevitauk

For the first few weeks after surgery, your initial results will be completely masked by swelling caused by tissue

/c/longevita

Even after the swelling goes down, it Be patient and don't worry, this is all part of the recovery process.

can take upwards of six months to see Please do not ever hesitate to contact your dedicated patient care specialist.